

Opinion

Environmentally Mediated Social Dilemmas

Sylvie Estrela,^{1,2,13,*} Eric Libby,^{3,4,13} Jeremy Van Cleve,^{5,13} Florence Débarre,⁶ Maxime Deforet,⁷ William R. Harcombe,⁸ Jorge Peña,^{9,10} Sam P. Brown,¹¹ and Michael E. Hochberg^{3,10,12,*}

By consuming and producing environmental resources, organisms inevitably change their habitats. The consequences of such environmental modifications can be detrimental or beneficial not only to the focal organism but also to other organisms sharing the same environment. Social evolution theory has been very influential in studying how social interactions mediated by public ‘goods’ or ‘bads’ evolve by emphasizing the role of spatial structure. The environmental dimensions driving these interactions, however, are typically abstracted away. We propose here a new, environment-mediated taxonomy of social behaviors where organisms are categorized by their production or consumption of environmental factors that can help or harm others in the environment. We discuss microbial examples of our classification and highlight the importance of environmental intermediates more generally.

Why the Environment Matters for Social Evolution

Organisms and their environment are inevitably coupled [1]. By consuming, transforming, and producing environmental resources, individuals modify their habitat. The consequences of this change can be damaging to a focal individual and its neighboring conspecifics, for example by producing harmful waste [2]. Conversely, environmental modification can be beneficial, for example when organisms invest in shelter or resource acquisition [3,4].

Social evolution (see [Glossary](#)) theory is an important framework for understanding behaviors that impact the fitness of other individuals, potentially resulting in (social) trait evolution. Social evolution theory has been very influential in the study of interactions among conspecifics, but it has traditionally overlooked the role of environmental dimensions in mediating such interactions and their impacts on the environment. The classical taxonomy of social evolution, first introduced by Hamilton [5–7], considers fitness-affecting interactions between focal and recipient individuals. ‘Helping’ interactions increase recipient fitness and can be either **altruistic** or **mutualistic** from the perspective of the focal actor, while ‘harming’ interactions decrease recipient fitness and can be either **selfish** or **spiteful** when they increase or decrease, respectively, the fitness of the actor [5]. The mechanisms of helping and harming often involve an environmental currency – for example a physical refuge or a chemical attack – but these environmental dimensions are typically abstracted away such that the amount of ‘help’ (or ‘harm’) is simply defined by a proxy such as the number of ‘helpers’ (or ‘harmers’) in the neighborhood of recipients [8,9].

The role of environmental intermediates in social interactions has received increasing attention in recent years owing to the growing realization that explicitly considering environmental dimensions is crucial for capturing realistic system complexity [1,8–12]. For example, if **cooperation** (or helping) among organisms improves the environment or prevents its degradation, then oscillations can emerge where cooperators and defectors alternately dominate as the environment cycles between replete and deplete states [11]. Thus, a more realistic and

Highlights

Organisms modify their environments in ways that can be beneficial or detrimental not only to themselves but also to others sharing the same environment.

Such niche-constructing or niche-destroying activities are often due to the production or consumption of environmental factors, such as resources, wastes, or toxins, which ultimately influence the ecology and evolution of social interactions.

We present a new, four-way classification of social behaviors where individual behaviors are categorized into producing/consuming an environmental factor, as well as into helping/harming others.

Although not immediately obvious, dispersal (the act of moving within a habitat or between habitats) is another mechanism by which organisms modify their environment.

An explicit representation of such environmentally mediated interactions is key to capturing realistic system complexity and can reveal some unexpected outcomes in social dilemmas.

¹Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT, USA

²Microbial Sciences Institute, Yale University, West Haven, CT, USA

³Santa Fe Institute, Santa Fe, NM, USA

⁴Department of Mathematics, Umeå University, Umeå, Sweden

⁵Department of Biology, University of Kentucky, Lexington, KY, USA

⁶Sorbonne Université, Université Pierre et Marie Curie (UPMC) Université Paris 06, Université Paris Diderot, Université Paris-Est Créteil, Centre National de la Recherche

complex theory of social evolution that includes these important feedbacks needs to account for **niche construction** effects [1].

At present, theoretical work on environmentally mediated social interactions has largely focused on simple nonspatial settings (except, notably, work on biofilms; e.g., [13–17]), while the broader field of social evolution has long recognized the importance of spatial structure and its role in generating genetic relatedness that can lead to the evolutionary success of cooperation and altruism (e.g., [5,18–20]). Mathematical models have shown the influence of spatial structure in determining the success of cooperative behaviors, particularly in the face of nonlinear group benefits (e.g., [21]), repeated interactions (e.g., [22,23]), or when individuals can move [24–28]. Nevertheless, these models often do not explicitly consider environmental dimensions. The few models that do consider explicit environmental variables suggest that, when cooperation improves the local environment, selection for cooperation can be very strong [29–32]. For example, Pepper and Smuts used an agent-based spatial model with conditional movement and environmental feedbacks to demonstrate that cooperation can be maintained in the presence of defection when cooperators improve the environment. Individuals are then less likely to leave locations with good environmental conditions [29].

In this Opinion we seek to address questions concerning how different types of social behavior can be maintained under the more realistic conditions of nonrandom (structured) interactions, explicit environmental currencies, and **dispersal** (or migration) of individuals (Figure 1). This is particularly relevant to cooperative behaviors, which have been shown to require one or more

Scientifique (CNRS) Institut de Recherche pour le Développement (IRD), Institut National de la Recherche Agronomique (INRA), Institute of Ecology and Environmental Sciences of Paris (iEES Paris), Paris, France

⁷Laboratoire Jean Perrin, CNRS, Sorbonne Université, Paris, France

⁸Department of Ecology, Evolution, and Behavior, University of Minnesota, Minneapolis, MN, USA

⁹Université Toulouse 1 Capitole, Toulouse, France


¹⁰Institute for Advanced Study in Toulouse, Toulouse, France

¹¹School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA, USA

¹²Institut des Sciences de l'Évolution, CNRS Unité Mixte de Recherche 5554, University of Montpellier, Montpellier, France

¹³These authors contributed equally

*Correspondence: sylvie.estrela@yale.edu (S. Estrela) and mhochber@univ-montp2.fr (M. Michael E.. Hochberg).



Trends in Ecology & Evolution

Figure 1. The Variety of Potential Interactions between Organisms and Their Environment. For simplicity, models of social evolution have often assumed that interactions between individuals are random, the physicochemical environment is implicit, and/or individuals cannot move within a patch or disperse between patches (top scenarios). We focus here on more realistic conditions, where interactions are nonrandom, the environmental dimension is explicit, and individuals can migrate (bottom scenarios), to investigate how these combined effects may influence the ecology and evolution of social behaviors. Note that we use the terms migration and dispersal interchangeably.

nonhomogeneous selective structures to persist (e.g., [23,31,33–36]). For instance, can ‘helper’ organisms sufficiently colocalize with the products of their helping activities to survive free-loaders? Can ‘harming’ organisms escape the environmental damage they create? Given the complexity of spatial systems of interacting organisms and environments, the answers to these and other related questions are far from straightforward, and require explicit mathematical models and controlled experiments (see Outstanding Questions).

Environmental Resources and the Tragedy of the Commons

Social dilemmas are loosely characterized by the conflict between the individual costs (or benefits) of particular actions and the benefits (or costs) of those actions when performed by enough members of a collective. In many cases, these actions involve the production or consumption of environmental resources that are valuable for the survival and reproduction of individuals in a group, where groups are defined by the ability of individual members to access those resources. These accessible resources are typically depletable, such as glucose concentration in a bacterial batch culture, and are referred to as ‘public goods’ or ‘common pool resources’. The social dilemma involved in maintaining public goods is often called the ‘tragedy of the commons’ [37]. The idea is that, if members of a social group use a shared limiting resource (‘the commons’) selfishly without restraint, then the commons will be depleted and eventually destroyed. Because organisms across the tree of life exhibit a variety of mechanisms involved in the production, acquisition, or maintenance of environmental resources and public goods, they all potentially face this dilemma [38,39]. Nonetheless, the degree to which evolution averts the tragedy of the commons is often a complex function of individual and collective interactions with the environment. We next present a classification of these interactions.

A Classification of Environmentally Mediated Social Behaviors

We present here a framework for classifying social dilemmas based on their environmental interactions that is inspired by Hamilton’s individual-based classification [5–7]. Our first step is to recognize that helping and harming behaviors can be classified as one of two distinct environmental interactions: (i) when the actor produces a helpful or harmful environmental factor, and (ii) when the actor consumes a harmful or helpful factor (Figure 2).

In our classification, a focal individual (actor) that produces a collectively beneficial good is referred to as an ‘enricher’, whereas an actor that degrades a damaging environmental factor is a ‘detoxifier’. Under the classical classification of social traits, both would be viewed as expressing a ‘helping’ behavior (with further categorization as either altruistic or mutualistic, depending on the degree to which the benefits return to the focal actor). Similarly, our new classification breaks down conventional ‘harming’ behaviors into ‘pollute’ and ‘deplete’, depending on whether the harming activity produces an environmental ‘bad’ or consumes a shared environmental ‘good’. Thus, what appears to be the same social behavior under the classical taxonomy of social evolution may actually be the result of opposing consume/produce strategies when the environment is explicitly considered. Therefore, to understand how social traits mediated through the environment evolve, it is crucial to explicitly consider the contrast between producing and consuming environmental goods and bads, and how this influences the outcome of social dilemmas at both temporal and spatial scales. We discuss below examples of our four-way classification with a focus on microbes.

Microbial Examples of the Proposed Classification

Microbes affect each other in many ways. These effects may arise through direct physical contact (e.g., attachment, toxin injection) or indirectly by simply sharing the same chemical

Glossary

Altruistic: a trait that increases the fitness of others (recipients) but decreases the fitness of the focal organism (actor).

Cheats: organisms that benefit from the costly behaviors of other organisms but without contributing, or contributing less.

Consumer–resource models: models that explicitly account for the dynamics of resources and of the organisms consuming those resources.

Cooperation: (or helping) social trait that increases the fitness of another individual (recipient).

Dispersal: a process by which one individual or a group of individuals move from one location to a new location. Dispersal can be active, through self-motility, or passive, as a result of external forces such as transportation through wind or water.

Lotka–Volterra models of

competition: a system of ordinary differential equations describing the dynamics of interacting populations (e.g., species). Common resources are not explicitly modeled, and the effect that one species has on itself and on others is implicitly modeled with interaction coefficients.

Mutualistic: a trait that increases the fitness of both recipient and actor.

Niche construction: a concept that broadly describes any organismal trait (behavior) that modifies its own environment and that of others. Such traits can have positive (constructive) or negative (destructive) effects on the environment.

Selfish: a trait that increases the fitness of the actor but decreases the fitness of the recipient.

Social evolution: an area of evolutionary biology that focuses on explaining the evolution of traits by a focal individual (actor) that have an effect (positive or negative) on the fitness of other individuals (recipients), and that evolved, in part, because of these effects.

Spiteful: a trait that decreases the fitness of both recipient and actor.




Figure 2. A Classification of Environmentally Mediated Social Behaviors. Individual behaviors are categorized into helping/harming others, and also into producing/consuming an environmental factor, yielding a four-way classification of behavior. Schematic summaries illustrate the signs of reciprocal organism (O)–environment (E) interactions under each scenario (a pointed arrow indicates a positive effect; a flat arrow indicates a negative effect). Effects on organisms are generally measured as changes in the density/amount of an environmental factor. For each scenario, all individuals in the population are assumed to have the same behavior, in other words they act and are affected by the environment in the same way. Thus, helping (or harming) others means helping (or harming) other individuals of the same type.

environment. Such environmental intermediates not only shape microbial composition ('who is there') but also the nature of their interaction ('what they do to each other'). We go beyond the traditional 'helping' versus 'harming' classification of social behavior. We discuss empirical and theoretical examples of our four-way classification – enrich, detoxify ('detox'), pollute, and deplete – to illustrate how considering explicit organism density and environmental dimensions can be important in understanding the ecology and evolution of social dilemmas. Although the mechanisms of enrich, detox, pollute, and deplete are discussed separately, they are clearly often intertwined and interdependent, contributing to the rich and complex diversity of microbe–microbe metabolic interactions in nature.

Enrich

Microbes can modify their environment by producing extracellular molecules that promote the growth of the focal individual and its neighboring conspecifics. Examples include exoenzymes that break down complex substrates into simple digestible compounds (e.g., glycoside hydrolases and proteases) [40], scavenging molecules that bind specific compounds in the environment and transport them back into the cell (e.g., iron-scavenging siderophores [41,42]), and structural compounds in biofilms that protect microbes from chemical or biological attacks [43–45], or place them near other important resources such as oxygen [46]. In some cases, such exoenzymes benefit the focal producer and conspecifics by directly harming others, such as antibiotics and toxins that kill or inhibit the growth of other interspecific organisms [47]. The production of such beneficial compounds is often costly and will only evolve if it sufficiently benefits the focal producer ('enricher') because otherwise it would be selected against and eventually lost from the population. Any benefits leaked to other neighboring organisms depend on how public or private these traits are.

Detoxify

A long-overlooked but increasingly important factor shaping microbial interactions is detoxification [48–58]. Examples include the production of enzymes such as β -lactamases that degrade β -lactam antibiotics [53,56], catalases that degrade hydrogen peroxide [51,58],

and metal-chelating siderophores that reduce toxic metal levels [54,55]. Detoxification not only benefits the focal producer ('detoxifier') but also protects neighboring cells living in the same environment. Because enzyme production is generally costly, detoxification is expected to have evolved to benefit the producer, and the benefits to others depend on the privatization level of detoxification [49,59]. Detoxification can nevertheless also be 'incidental' (i.e., is not selected for), such as when an organism feeds on compounds that are toxic to others but not to itself [48,60].

Pollute

Microbial cells commonly excrete metabolic waste products such as lactic acid, acetate, and hydrogen. These metabolic byproducts may become toxic, especially at high concentrations, suppressing the growth of not only the producer cell but also of any nearby susceptible cells (e.g., by changing the pH [61]). In some cases, however, such metabolic waste can be a resource for another species, allowing both polluters (waste producers) and detoxifiers (cross-feeders) to potentially benefit from the interaction if the mutual benefits of food for detoxification outweigh the costs of competition for shared resources [48]. Such syntrophic interactions are common between hydrogen-producing bacteria and methanogens. While the build-up of hydrogen inhibits the growth of the producing bacteria, methanogens are able to use hydrogen for growth, and thus keep hydrogen levels low, allowing their hydrogen-producing partner to grow [62,63].

Deplete


Competition for resources – occurring when organisms feed on shared, limiting resources (also known as 'exploitation competition') – is a key factor shaping microbial interactions [64]. Although resource competition is often initially weak, competition increases as cells grow and resources become depleted. In an unstructured environment with a single resource, theory predicts that the more-competitive strain displaces the other strains, thus resulting in competitive exclusion [65]. However, the presence of trade-offs [66,67], such as 'fast-growing but low-yield' versus 'slow-growing but high-yield', can promote species coexistence even when organisms compete for a single resource. From a social evolution perspective, a slow-growing and high-yield strategy can be viewed as a form of cooperation because resources are used in a more prudent way compared to a fast-growing and low-yield strategy. Thus, the evolutionary dilemma is how to maintain prudent (cooperative) resource use in the face of rapacious (noncooperative) individuals that eventually deplete the commons. A solution to this dilemma is spatial structure that enables cooperators to preferentially interact with other cooperators and remain protected from the local effects of noncooperators [13,68–70].

The Value of Explicit Environmental Variables

Models of resource competition have often assumed that resources are either implicit (e.g., **Lotka–Volterra models of competition** [71]) or explicit, but with a single resource pool (e.g., classic **consumer–resource models** with a single limiting resource [72]). Microbial cells, however, are often surrounded by a variety of genotypes with different resource-use strategies and diverse resource pools (supplied exogenously and/or as byproducts of cell metabolism). Classic population ecology models or resource-based models with a single resource can therefore lead to an oversimplified depiction of organism–organism interactions and, in some cases, fail to predict the dynamics of microbial interactions [73,74]. Work combining these and other explicit metabolic models with spatially explicit environmental models [75–77] have recently received more attention, and promise to yield a wealth of insights into more realistic representations of ecological interactions in nature.

Environment-Mediated Organism–Organism interactions

If we abstract away the environment and consider two organisms and the various positive and negative interactions they can exert on themselves and each other, we find 10 possible scenarios after removing symmetries (Figure 3). Six of these interaction scenarios can occur via a single environmental compound (A, D–G, J). Interestingly, some reduced (environmentally implicit) systems map to two different environmentally mediated social interactions (A, D, E, J), while others map to only one (F, G). The other four scenarios, however, are impossible to achieve with only one environmental intermediate (B, C, H, I). To see this, consider scenario B in which one organism (O_1) positively affects itself and the other organism (O_2). This could occur



Trends in Ecology & Evolution

Figure 3. Environment-Mediated Organism–Organism Interactions. Illustrated are the possible combinations of positive (red) and negative (blue) interaction scenarios between two organisms (O_1 and O_2) of different types when the environment is implicit, taking into account the effect of each organism on themselves and individuals of the same type (illustrated as an arrow end pointing towards self), as well as the effect on others (pointing towards the other type). Note that here we remove symmetries: that is, when species O_1 and O_2 are interchangeable. Six of the reduced systems can be used to describe all 10 unique scenarios involving a single environmental intermediate (E) and combinations of the environmental modifications enrich, detox, pollute, deplete (A, D–G, J). The other four reduced systems cannot be described with a single environmental factor and at least two are needed (B, C, H, I). Note that we only represent environmental variables that directly mediate the interaction between the two organisms. Environmental factors that affect the growth of one organism but not the other are omitted.

through the removal of a toxic product (detox) or the production of a beneficial product (enrich). In addition, organism O_2 interacts positively with O_1 but negatively with itself (e.g., by producing a waste product that is toxic to itself but on which O_1 can feed, or by feeding on a compound that is toxic to O_1). A simple ' O_1 - E_1 - O_2 ' schematic is limited because it cannot capture the possibility that the single environmental compound (E_1) that helped both O_1 and O_2 can also help O_1 and hurt O_2 . Therefore, we need a second environmental factor (E_2) to be able to capture such scenarios.

Furthermore, the mappings from the reduced system to the environmentally explicit system are not always identical. For instance, scenario J in the reduced system corresponds to a case where each organism negatively affects itself and the other organism. In the environmentally explicit model, this can be either deplete/deplete or pollute/pollute. In the case of deplete/deplete, both organisms consume the same resource, and hence negatively affect themselves and each other by removing the resource. In the case of pollute/pollute, both organisms produce the same waste product that negatively affects everyone. Importantly, these two scenarios produce different dynamical models, especially regarding the dynamics of E . It is also worth noting that some implicit models where O_1 and O_2 have the same strategy lead to explicit models where O_1 and O_2 also have the same strategy (J), but in other cases O_1 and O_2 must have different strategies (F, G).

Spatial and Temporal Dynamics of Environmentally Mediated Social Dilemmas

Although it is well known in social evolution theory that the spatial scale at which cooperation and conflict over resources occurs can have a large impact on evolved levels of cooperation [78–81], it is possible that unexpected social evolutionary effects emerge as a consequence of interactions between environmental, spatial, and temporal parameters. To illustrate these possibilities, we describe below how the viscosity (spatial scale) and persistence (temporal scale) of environmental factors can affect the evolution of cooperative traits and, in turn, the spatial arrangement of cooperators and noncooperators.

Environmental Viscosity: Cell Dispersal and Exoproduct Diffusion

Environmental mediation allows social interactions to take place at larger spatial scales via diffusible compounds. Whether or not spatial dynamics favor enriching strategies depends on environmental viscosity and its effect on the levels of cell dispersal and public goods diffusion [80,82]. In the case of high viscosity, where both cell dispersal and public goods diffusion are limited, the benefits of public goods can be channeled between adjacent and likely related organisms. This acts to privatize the produced goods and can promote cooperation (e.g., [82,83]). However, if viscosity is such that cell dispersal is limited while public goods diffusion remains high, then the spatial scale over which cooperative interactions occur becomes larger than the spatial scale of competition [84], and viscosity will not favor the evolution of cooperation (e.g., [85]).

The interplay of social interactions and environmental viscosity has received particular attention in the study of microbial biofilms [17]. To capture realistic features of microbial biofilms as well as the chemical/physical properties of the environment, spatially and mechanistically explicit individual-based models have been shown to be a powerful approach [15]. Such models can provide important insights into how factors such as exoproduct diffusion, population growth, and social interactions affect the eco-evolutionary dynamics of these spatially-structured systems [17,86]. For instance, the extracellular matrix that generates biofilms was initially thought of as a public good that could be exploited by nonproducers. However, individual-

based models that incorporated nutrient diffusion and cell-limited dispersal demonstrated that polymer secretion is often a competitive strategy that lifts producer genotypes into favorable environments while suffocating nonproducer genotypes [14]. Further, individual-based models have revealed how social interactions shape the arrangement of genotypes, and therefore selection, in structured systems. While cells with cooperative genotypes will mix as they grow and divide, competing genotypes become segregated as a result of growth [87,88]. This can lead to social insulation of cooperative genotypes from **cheating** mutants, and enhance selection for cooperation [16,89].

Environmental Persistence: Organism Lifespan and Exoproduct Durability

Another important consequence of environmental mediation is that, when modified environmental factors persist for long enough, their effects can be transmitted across generations [8,10] (such that there is ecological inheritance, *sensu* [1,90]). This means that actors interact not only with existing relatives but also with relatives living in the future who are not in direct competition with actors in the present [91,92]. This decouples kin-selected benefits and costs resulting from kin competition and can increase selection pressure on helping traits, including altruistic behaviors that would be selected against when actors and recipients only interact during the same generation [91]. For durable public goods that are nonrival (i.e., consumption does not reduce availability to others), investments by earlier generations can lead to loss of cooperative genotypes in subsequent generations – unless cooperative behaviors are under regulatory control [8,10]. For microbial interactions, these observations suggest that not only the viscosity of the medium is important but also the durability of secreted compounds relative to the lifespan of organisms. In the case where exoproduct durability is sufficiently larger than the lifespan of the focal producer, ecological inheritance can operate, thus paving the way for posthumous niche construction [92].

Environmentally Mediated Conditional Dispersal

The production and consumption of environmental factors (as in the enrich, detox, pollute, deplete scenarios) are one of two ways that organisms can engage in niche-constructing activities. The other way is active movement of organisms within a patch and dispersal between patches. By dispersing, an individual necessarily changes the patch from which it dispersed, as well as the patch to which it migrates [1]. One inclusive fitness benefit of dispersal is the ability of an individual to avoid competing for resources with its relatives in its natal patch. Although this topic has received considerable theoretical attention [27,93–100], previous work mostly considers the environment implicitly and does not detail the helpful or harmful environmental factors that affect survival and reproduction (i.e., our classification above). One crucial consequence of omitting these environmental factors is the inability to consider how dispersal can evolve to be conditional on the state of the local environment [101], which is itself affected by organismal niche construction. Because many social interactions can be produced by multiple explicit interactions with the environment (Figure 3), the effect of dispersal conditional on environmental state is likely to depend on these explicit interactions.


To see how dispersal conditional on environmental variables may be important, consider scenario J in Figure 3, whose implicit representation consists of negative effects of each organism on itself and on the other organism. Scenario J can be represented by two explicit environmental interactions: (i) both organisms negatively affect the environmental compound which in turn positively affects them, as occurs in a ‘deplete’ scenario, or (ii) both organisms positively affect the environment which negatively affects them, as occurs in a ‘pollute’ scenario. Organismal growth rates increase with greater concentrations of the environmental compound in the ‘deplete’ scenario, and decrease in the ‘pollute’ scenario. Thus, all else equal,

dispersal rates away from the current patch may evolve to be higher for low concentrations of the environmental product in the 'deplete' scenario but lower for high concentrations of the product in the 'pollute' scenario (Box 1). Thus, a single implicit social interaction can generate different dispersal mechanisms depending on the explicit environmental interactions. A more complex scenario is presented in Box 1.

Further, dispersal conditional on explicit environmental variables may affect the conditions under which different types of social behaviors evolve. For example, models have shown that cooperative interactions that increase group survival or carrying capacity can easily evolve [102,103]. In effect, these cases represent a type of 'enrich' scenario whereby cooperators create larger and more persistent groups. However, if dispersal is allowed to be conditional on local resource concentrations, then noncooperators are no longer so easily trapped in unproductive groups that become extinct and might preferentially disperse from low-resource patches to high-resource patches with more cooperators. Such conditional dispersal would likely erode the likelihood that cooperators enrich the environment for other cooperators and diminish selection for cooperation in these scenarios. Interestingly, there might also be selection for conditional dispersal of cooperators away from low-resource patches to high-resource patches. Because high resource concentrations typically cannot be generated by groups composed mostly of noncooperators, they might serve as an environmental cue of the presence of other cooperators, and could create the type of positive assortment among cooperators that is necessary to maintain them in the presence of noncooperators [23,31]. In fact, Pepper and Smuts [29] studied this type of conditional dispersal and found that it can generate significant assortment between cooperators. More generally, conditional dispersal might either enhance or erode cooperation, depending on the details of the environmental interactions. Thus, not including explicit environmental interactions could exclude important ecological and evolutionary dynamics in social dilemmas.

Box 1. The Value of Explicit Environment Representation in the Evolution of Dispersal

The interplay between dispersal and environmentally mediated social dilemmas can benefit from explicit representations of the environment in theoretical models. For example, consider scenario F in Figure 3. In an implicit representation of the environment, organisms O_1 and O_2 appear to be identical in kind: both help themselves and harm the other. Now suppose that each organism can disperse to a new environment. Under the implicit representation, there is no suggestion that O_1 and O_2 would follow different dispersal strategies. Nonetheless, an explicit representation of the environment shows that they are affected by the environment differently: one enriches the environment while the other detoxifies it. Thus, we would expect each organism to follow a different dispersal strategy as a function of environmental state: the enrich organism has a higher growth rate in higher concentrations of E, and as such should remain in high-E patches (lower dispersal rate from such patches), while the detox organism is harmed by higher concentrations of E and so should move away from high-E patches (high dispersal rate) (Figure 1A). In a sense, this agrees with the implicit environmental model in that the two organisms should disperse away from one another because they harm each other. Yet, the explicit environmental model suggests a mechanism by which a patchy environment of high/low concentrations of E might facilitate a corresponding spatial assortment of the organisms. Let us now turn to scenario J in Figure 3. Again, in an implicit representation of the environment, organisms O_1 and O_2 appear to be identical in kind: both harm themselves and harm each other. In this case, however, the two organisms interact with the environment in the same way and two different environmentally mediated interactions are possible: either deplete/deplete or pollute/pollute. We therefore expect each organism to adopt the same dispersal strategy. Importantly, however, the specific strategy they adopt – a higher or lower dispersal rate – depends on how the explicit environment is represented (Figure 1B). Such a contrast would be difficult to capture with an environmentally implicit model. This is also the case in scenario E in Figure 3, in which the two organisms interact with the environment in a different way, and therefore should adopt opposite dispersal strategies. Nevertheless, whether each organism should stay or leave a patch with a particular concentration of E is determined by how the environment is represented (Figure 1C). Together, these examples highlight how considering both environmental and social conditions explicitly can provide important insights for understanding and predicting the evolution of conditional dispersal.



Trends in Ecology & Evolution

Figure 1. The Value of Explicit Environment Representation for Predicting the Evolution of Dispersal. Plots show the predicted growth rates and dispersal probabilities of two organisms (O_1 and O_2) interacting through a shared environmental factor (E) for three of the scenarios represented in Figure 3. The lines are intended to represent the direction of the trend (increasing/decreasing) and not the form (depicted as linear, for simplicity). Here the dispersal probability of O_1 (or O_2) means the probability that an organism will move away from a patch with density E .

Concluding Remarks

The typology we describe in Figure 2 aims to catalyze a better mechanistic understanding of the ecological and evolutionary dynamics of environmentally mediated social dilemmas. Nevertheless, the simple typology of social dilemmas initiated by Hamilton that classifies behaviors strictly based on their fitness effects is still an important conceptual tool [5]. It produces many insights including the role of genetic identity and coancestry in the evolution of altruism [5], as well as the effect of local competition for resources and other demographic forces in altering selection for or against cooperation [18,19,102]. Therefore, a crucial task is to develop formal mathematical connections between the environmentally mediated and classical social dilemma typologies.

Under some circumstances, environmentally implicit approaches will accurately represent social dilemmas and their dynamics. Evidently, this is increasingly the case when habitats become unstructured, when social interactions are more direct and less mediated by elements of the environment (as occurs with mechanisms requiring direct cell-to-cell contact, e.g., competitor killing via the type VI secretion systems, T6SS) [104], and when the environmental dimension is coupled to producers (e.g., when there is fast turnover of the environmental currency and/or a short lag between changes in focal producer and public good/bad densities [8]). However, we suspect that such simplicity, even if common, is not representative of many environments. Thus, a more explicit framework such as that presented here is necessary for structured environments and environmentally mediated social interactions. Attaining both generality and accuracy in such a framework will be challenging given the stunning diversity

of social systems [105] and their complex associations with the environment [1]. Nevertheless, the ubiquitous importance of social interactions and how dilemmas are mediated by ecology and evolution argue for an environmentally inclusive framework and future investigation using mathematical, behavioral, molecular, and genomic tools. Although our examples focus on microbial systems, we believe that our four-way classification will also be useful for understanding interactions mediated through the environment in other systems such as social animals and humans.

Acknowledgments

We wish to thank the Santa Fe Institute for hosting two working groups in 2016 and 2017, during which this research was conducted, and four anonymous reviewers for helpful suggestions. F.D. received funding from Agence Nationale de la Recherche (ANR, grant 14-ACHN-0003). J.P. gratefully acknowledges financial support from the ANR-Labex Institute for Advanced Study in Toulouse (IAST). M.E.H. thanks the McDonnell Foundation (Studying Complex Systems research award 220020294).

References

- Odling-Smee, F.J. *et al.* (2003) *Niche Construction: The Neglected Process in Evolution*, Princeton University Press
- Bédhomme, S. *et al.* (2005) Pollution by conspecifics as a component of intraspecific competition among *Aedes aegypti* larvae. *Ecol. Entomol.* 30, 1–7
- Crespi, B.J. (2001) The evolution of social behavior in microorganisms. *Trends Ecol. Evol.* 16, 178–183
- West, S.A. *et al.* (2006) Social evolution theory for microorganisms. *Nat. Rev. Microbiol.* 4, 597–607
- Hamilton, W.D. (1964) The genetical evolution of social behaviour. I. *J. Theor. Biol.* 7, 1–16
- Hamilton, W.D. (1964) The genetical evolution of social behaviour. II. *J. Theor. Biol.* 7, 17–52
- Hamilton, W.D. (1970) Selfish and spiteful behaviour in an evolutionary model. *Nature* 228, 1218–1220
- Brown, S.P. and Taddei, F. (2007) The durability of public goods changes the dynamics and nature of social dilemmas. *PLoS One* 2, e593
- McNally, L. and Brown, S.P. (2015) Building the microbiome in health and disease: niche construction and social conflict in bacteria. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 370, 20140298
- Kümmerli, R. and Brown, S.P. (2010) Molecular and regulatory properties of a public good shape the evolution of cooperation. *Proc. Natl. Acad. Sci. U. S. A.* 107, 18921–18926
- Weitz, J.S. *et al.* (2016) An oscillating tragedy of the commons in replicator dynamics with game–environment feedback. *Proc. Natl. Acad. Sci. U. S. A.* 113, E7518–E7525
- Benincà, E. *et al.* (2015) Species fluctuations sustained by a cyclic succession at the edge of chaos. *Proc. Natl. Acad. Sci. U. S. A.* 112, 6389–6394
- Kreft, J.U. (2004) Biofilms promote altruism. *Microbiology* 150, 2751–2760
- Xavier, J.B. and Foster, K.R. (2007) Cooperation and conflict in microbial biofilms. *Proc. Natl. Acad. Sci. U. S. A.* 104, 876–881
- Nadell, C.D. *et al.* (2010) Emergence of spatial structure in cell groups and the evolution of cooperation. *PLoS Comput. Biol.* 6, e1000716
- Mitri, S. *et al.* (2011) Social evolution in multispecies biofilms. *Proc. Natl. Acad. Sci. U. S. A.* 108, 10839–10846
- Nadell, C.D. *et al.* (2016) Spatial structure, cooperation and competition in biofilms. *Nat. Rev. Microbiol.* 14, 589–600
- Lehmann, L. and Rousset, F. (2010) How life history and demography promote or inhibit the evolution of helping behaviours. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 365, 2599–2617
- Taylor, P.D. (1992) Altruism in viscous populations – an inclusive fitness model. *Evol. Ecol.* 6, 352–356
- Lehmann, L. *et al.* (2007) The evolution of helping and harming on graphs: the return of the inclusive fitness effect. *J. Evol. Biol.* 20, 2284–2295
- Peña, J. *et al.* (2015) Evolutionary dynamics of collective action in spatially structured populations. *J. Theor. Biol.* 382, 122–136
- Akçay, E. and van Cleve, J. (2012) Behavioral responses in structured populations pave the way to group optimality. *Am. Nat.* 179, 257–269
- van Cleve, J. and Akçay, E. (2014) Pathways to social evolution: reciprocity, relatedness, and synergy. *Evolution* 68, 2245–2258
- Hauert, C. and Doebeli, M. (2004) Spatial structure often inhibits the evolution of cooperation in the snowdrift game. *Nature* 428, 643–646
- Aktipis, C.A. (2004) Know when to walk away: contingent movement and the evolution of cooperation. *J. Theor. Biol.* 231, 249–260
- Aktipis, C.A. (2011) Is cooperation viable in mobile organisms? Simple walk away rule favors the evolution of cooperation in groups. *Evol. Hum. Behav.* 32, 263–276
- Hochberg, M.E. *et al.* (2008) The coevolution of cooperation and dispersal in social groups and its implications for the emergence of multicellularity. *BMC Evol. Biol.* 8, 238
- Parvinen, K. (2013) Joint evolution of altruistic cooperation and dispersal in a metapopulation of small local populations. *Theor. Popul. Biol.* 85, 12–19
- Pepper, J.W. and Smuts, B.B. (2002) A mechanism for the evolution of altruism among nonkin: positive assortment through environmental feedback. *Am. Nat.* 160, 205–213
- Pepper, J.W. (2007) Simple models of assortment through environmental feedback. *Artif. Life* 13, 1–9
- Lehmann, L. and Keller, L. (2006) The evolution of cooperation and altruism – a general framework and a classification of models. *J. Evol. Biol.* 19, 1365–1376
- Kéfi, S. *et al.* (2008) Evolution of local facilitation in arid ecosystems. *Am. Nat.* 172, E1–E17
- West, S.A. *et al.* (2007) Evolutionary explanations for cooperation. *Curr. Biol.* 17, R661–R672
- Nowak, M.A. (2006) Five rules for the evolution of cooperation. *Science* 314, 1560–1563
- Fletcher, J.A. and Doebeli, M. (2009) A simple and general explanation for the evolution of altruism. *Proc. Biol. Sci.* 276, 13–19
- Frank, S.A. (1994) Genetics of mutualism: the evolution of altruism between species. *J. Theor. Biol.* 170, 393–400
- Hardin, G. (1968) The tragedy of the commons. *Science* 162, 1243–1248
- Rankin, D.J. *et al.* (2007) The tragedy of the commons in evolutionary biology. *Trends Ecol. Evol.* 22, 643–651

Outstanding Questions

To what extent do environmental, metabolic, and spatial parameters impact the evolution of enrich, detox, pollute, or deplete interaction scenarios, and what is their relative importance?

Under which mechanistic and environmental conditions do implicit and explicit models generate different qualitative predictions? When does abstracting away the environment make reasonable predictions?

How does the interplay between dispersal and environmentally mediated interactions influence the evolutionary dynamics of social dilemmas and dispersal? What can we learn from using more environmentally explicit models of dispersal evolution?

Less is known about organisms that harm and about how the environment mediates these interactions. Can ‘harming’ organisms escape the environmental damage they create?

What is the role of positive versus negative ecological effects in describing cheating strategies whereby organisms profit from the cooperative traits of others without contributing themselves?

How do temporally and spatially dynamic environments influence the tragedy of the commons?

39. Levin, S.A. (2014) Public goods in relation to competition, cooperation, and spite. *Proc. Natl. Acad. Sci. U. S. A.* 111, 10838–10845
40. Rakoff-Nahoum, S. *et al.* (2016) The evolution of cooperation within the gut microbiota. *Nature* 533, 255–259
41. Griffin, A.S. *et al.* (2004) Cooperation and competition in pathogenic bacteria. *Nature* 430, 1024–1027
42. Cordero, O.X. *et al.* (2012) Public good dynamics drive evolution of iron acquisition strategies in natural bacterioplankton populations. *Proc. Natl. Acad. Sci. U. S. A.* 109, 20059–20064
43. Nadell, C.D. *et al.* (2015) Extracellular matrix structure governs invasion resistance in bacterial biofilms. *ISME J.* 9, 1700–1709
44. Vidakovic, L. *et al.* (2018) Dynamic biofilm architecture confers individual and collective mechanisms of viral protection. *Nat. Microbiol.* 3, 26–31
45. Kovacs, A.T. *et al.* (2012) The protective layer of biofilm: a repellent function for a new class of amphiphilic proteins. *Mol. Microbiol.* 85, 8–11
46. Rainey, P.B. and Rainey, K. (2003) Evolution of cooperation and conflict in experimental bacterial populations. *Nature* 425, 72–74
47. Riley, M.A. *et al.* (1999) The ecological role of bacteriocins in bacterial competition. *Trends Microbiol.* 7, 129–133
48. Estrela, S. *et al.* (2012) From metabolism to ecology: cross-feeding interactions shape the balance between polymicrobial conflict and mutualism. *Am. Nat.* 180, 566–576
49. Estrela, S. *et al.* (2016) Private benefits and metabolic conflicts shape the emergence of microbial interdependencies. *Environ. Microbiol.* 18, 1415–1427
50. Yurtsev, E.A. *et al.* (2016) Oscillatory dynamics in a bacterial cross-protection mutualism. *Proc. Natl. Acad. Sci. U. S. A.* 113, 6236–6241
51. Morris, J.J. *et al.* (2011) Dependence of the cyanobacterium *Prochlorococcus* on hydrogen peroxide scavenging microbes for growth at the ocean's surface. *PLoS One* 6, e16805
52. Sorg, R.A. *et al.* (2016) Collective resistance in microbial communities by intracellular antibiotic deactivation. *PLoS Biol.* 14, e2000631
53. Medaney, F. *et al.* (2016) Live to cheat another day: bacterial dormancy facilitates the social exploitation of β -lactamases. *ISME J.* 10, 778–787
54. O'Brien, S. *et al.* (2018) No effect of intraspecific relatedness on public goods cooperation in a complex community. *Evolution* 72, 1165–1173
55. O'Brien, S. *et al.* (2014) Social evolution of toxic metal bioremediation in *Pseudomonas aeruginosa*. *Proc. R. Soc. B Biol. Sci.* 281, 20140858
56. Frost, I. *et al.* (2018) Cooperation, competition and antibiotic resistance in bacterial colonies. *ISME J.* 12, 1582–1593
57. Estrela, S. and Brown, S.P. (2018) Community interactions and spatial structure shape selection on antibiotic resistant lineages. *PLoS Comput. Biol.* 14, e1006179
58. Morris, J.J. *et al.* (2014) Coexistence of evolving bacteria stabilized by a shared black queen function. *Evolution* 68, 2960–2971
59. Morris, J.J. (2015) Black Queen evolution: the role of leakiness in structuring microbial communities. *Trends Genet.* 31, 475–482
60. Cowan, S.E. *et al.* (2000) Commensal interactions in a dual-species biofilm exposed to mixed organic compounds. *Appl. Environ. Microbiol.* 66, 4481–4485
61. Ratzke, C. and Gore, J. (2018) Modifying and reacting to the environmental pH can drive bacterial interactions. *PLoS Biol.* 16, e2004248
62. Morris, B.E.L. *et al.* (2013) Microbial syntrophy: interaction for the common good. *FEMS Microbiol. Rev.* 37, 384–406
63. Hillesland, K.L. and Stahl, D.A. (2010) Rapid evolution of stability and productivity at the origin of a microbial mutualism. *Proc. Natl. Acad. Sci. U. S. A.* 107, 2124–2129
64. Ghoul, M. and Mitri, S. (2016) The ecology and evolution of microbial competition. *Trends Microbiol.* 24, 833–845
65. Hardin, G. (1960) The competitive exclusion principle. *Science* 131, 1292–1297
66. Bohannan, B.J.M. *et al.* (2002) Trade-offs and coexistence in microbial microcosms. *Antonie Van Leeuwenhoek* 81, 107–115
67. Litchman, E. *et al.* (2015) Microbial resource utilization traits and trade-offs: implications for community structure, functioning, and biogeochemical impacts at present and in the future. *Front. Microbiol.* 6, 65
68. Pfeiffer, T. *et al.* (2001) Cooperation and competition in the evolution of ATP-producing pathways. *Science* 292, 504–507
69. Kerr, B. *et al.* (2006) Local migration promotes competitive restraint in a host–pathogen 'tragedy of the commons'. *Nature* 442, 75–78
70. Nahum, J.R. *et al.* (2011) Evolution of restraint in a structured rock–paper–scissors community. *Proc. Natl. Acad. Sci. U. S. A.* 108, 10831–10838
71. Volterra, V. (1928) Variations and fluctuations of the number of individuals in animal species living together. *ICES J. Mar. Sci.* 3, 3–51
72. Chesson, P. (1990) MacArthur's consumer–resource model. *Theor. Popul. Biol.* 37, 26–38
73. Momeni, B. *et al.* (2017) Lotka–Volterra pairwise modeling fails to capture diverse pairwise microbial interactions. *eLife* 6, 205
74. Goldford, J.E. *et al.* (2018) Emergent simplicity in microbial community assembly. *Science* 361, 469–474
75. Harcombe, W.R. *et al.* (2014) Metabolic resource allocation in individual microbes determines ecosystem interactions and spatial dynamics. *Cell Rep.* 7, 1104–1115
76. Chacón, J.M. *et al.* (2018) The spatial and metabolic basis of colony size variation. *ISME J.* 12, 669–680
77. Xavier, J.B. *et al.* (2005) A framework for multidimensional modelling of activity and structure of multispecies biofilms. *Environ. Microbiol.* 7, 1085–1103
78. Allison, S.D. (2005) Cheaters, diffusion and nutrients constrain decomposition by microbial enzymes in spatially structured environments. *Ecol. Lett.* 8, 626–635
79. Driscoll, W.W. and Pepper, J.W. (2010) Theory for the evolution of diffusible external goods. *Evolution* 64, 2682–2687
80. Allen, B. *et al.* (2013) Spatial dilemmas of diffusible public goods. *eLife* 2, e01169
81. Dobay, A. *et al.* (2014) Interaction effects of cell diffusion, cell density and public goods properties on the evolution of cooperation in digital microbes. *J. Evol. Biol.* 27, 1869–1877
82. Kümmerli, R. *et al.* (2014) Habitat structure and the evolution of diffusible siderophores in bacteria. *Ecol. Lett.* 17, 1536–1544
83. Kümmerli, R. *et al.* (2009) Viscous medium promotes cooperation in the pathogenic bacterium *Pseudomonas aeruginosa*. *Proc. Biol. Sci.* 276, 3531–3538
84. Platt, T.G. and Bever, J.D. (2009) Kin competition and the evolution of cooperation. *Trends Ecol. Evol.* 24, 370–377
85. Le Gac, M. and Doebeli, M. (2010) Environmental viscosity does not affect the evolution of cooperation during experimental evolution of colicigenic bacteria. *Evolution* 64, 522–533
86. Steenackers, H.P. *et al.* (2016) Experimental evolution in biofilm populations. *FEMS Microbiol. Rev.* 40, 373–397
87. Estrela, S. and Brown, S.P. (2013) Metabolic and demographic feedbacks shape the emergent spatial structure and function of microbial communities. *PLoS Comput. Biol.* 9, e1003398
88. Momeni, B. *et al.* (2013) Strong inter-population cooperation leads to partner intermixing in microbial communities. *eLife* 2, e00230
89. Harcombe, W.R. *et al.* (2016) Adding biotic complexity alters the metabolic benefits of mutualism. *Evolution* 70, 1871–1881
90. Odling-Smee, F.J. *et al.* (1996) Niche construction. *Am. Nat.* 147, 641–648

91. Lehmann, L. (2007) The evolution of trans-generational altruism: kin selection meets niche construction. *J. Evol. Biol.* 20, 181–189
92. Lehmann, L. (2008) The adaptive dynamics of niche constructing traits in spatially subdivided populations: evolving posthumous extended phenotypes. *Evolution* 62, 549–566
93. Hamilton, W.D. and May, R.M. (1977) Dispersal in stable habitats. *Nature* 269, 578–581
94. Taylor, P.D. and Frank, S.A. (1996) How to make a kin selection model. *J. Theor. Biol.* 180, 27–37
95. Gandon, S. and Michalakis, Y. (1999) Evolutionarily stable dispersal rate in a metapopulation with extinctions and kin competition. *J. Theor. Biol.* 199, 275–290
96. Perrin, N. and Lehmann, L. (2001) Is sociality driven by the costs of dispersal or the benefits of philopatry? A role for kin-discrimination mechanisms. *Am. Nat.* 158, 471–483
97. Le Galliard, J.F. *et al.* (2005) Adaptive evolution of social traits: origin, trajectories, and correlations of altruism and mobility. *Am. Nat.* 165, 206–224
98. Mouden El, C. and Gardner, A. (2008) Nice natives and mean migrants: the evolution of dispersal-dependent social behaviour in viscous populations. *J. Evol. Biol.* 21, 1480–1491
99. Lehmann, L. and Perrin, N. (2002) Altruism, dispersal, and phenotype-matching kin recognition. *Am. Nat.* 159, 451–468
100. Mullon, C. *et al.* (2018) Social polymorphism is favoured by the co-evolution of dispersal with social behaviour. *Nat. Ecol. Evol.* 2, 132–140
101. Rodrigues, A.M.M. and Johnstone, R.A. (2014) Evolution of positive and negative density-dependent dispersal. *Proc. Biol. Sci.* 281, 20141226
102. Lehmann, L. *et al.* (2006) Population demography and the evolution of helping behaviors. *Evolution* 60, 1137–1151
103. Powers, S.T. and Lehmann, L. (2013) The co-evolution of social institutions, demography, and large-scale human cooperation. *Ecol. Lett.* 16, 1356–1364
104. McNally, L. *et al.* (2017) Killing by type VI secretion drives genetic phase separation and correlates with increased cooperation. *Nat. Commun.* 8, 14371
105. Bourke, A.F.G. (2011) *Principles of Social Evolution*, Oxford University Press